

## Wellness Within Your Reach

1st Floor, ADOL House, 15 CIPM Road, Alausa, Ikeja, Lagos. TEL: +234 704 770 0006 Email: hello@josmolgroup.com

Patient Name: Referred by:

Age/Sex : Registration Date :

Lab ID No : Collected Date :

Clinical Summary:

## KIDNEY FUNCTION TEST

TEST RESULT UNIT REFERENCE

2

**Comment:** GFR Category (ml/min/1.73m²)

G1 :  $\geq 90$  Normal or High G2 : 60 - 89 Mildly decreased

G3a : 45 - 59 Mildly to moderately decreased G3b : 30 - 44 Moderately to severely decreased

G4 : 15 - 29 Severely decreased G5 : ≤ 15 Kidney failure

Note that analytical interference may sometimes cause erroneous results. Please correlate all laboratory results with your clinician

Dr. C.O. Famuyiwa

FMCPath Consultant Pathologist



## Wellness Within Your Reach

1st Floor, ADOL House, 15 CIPM Road, Alausa, Ikeja, Lagos. TEL: +234 704 770 0006 Email: hello@josmolgroup.com

Patient Name: Referred by:

Age/Sex : Registration Date :

Lab ID No : Collected Date :

Clinical Summary:

## PROSTATE SPECIFIC ANTIGEN (PSA)

TEST RESULT UNIT REFERENCE

An elevated PSA level is suggestive of cancer but there are other possible causes of an elevated PSA level. These include benign conditions such as Prostatis and Benign Prostatic Hyperplasia (BPH). Also, a recent rectal examination can cause a rise in PSA levels. An elevated PSA level done alone does not give enough information to distinguish between benign prostatic conditions and cancer. A prostate biopsy is the confirmatory test to diagnose cancer of the prostate

Note that analytical interference may sometimes cause erroneous results. Please correlate all laboratory results with your clinician

Dr. C.O. Famuyiwa

FMCPath Consultant Pathologist

